



Long-term survival and major outcomes in post-cardiotomy extracorporeal membrane oxygenation for adult patients in cardiogenic shock

Paolo Meani^{1,2}, Matteo Matteucci^{1,3}, Federica Jiritano¹, Dario Fina¹, Francesco Panzeri⁴, Giuseppe M. Raffa^{1,5,6}, Mariusz Kowalewski¹, Nuccia Morici^{7,8}, Giovanna Viola⁷, Alice Sacco⁷, Fabrizio Oliva⁷, Amal Alyousif⁴, Sam Heuts¹, Martijn Gilbers¹, Rick Schreurs¹, Jos Maessen¹, Roberto Lorusso¹

¹Department of Cardio-Thoracic Surgery, Maastricht University Medical Centre (MUMC+), Maastricht, The Netherlands; ²Department of Intensive Care Unit, Maastricht University Medical Centre (MUMC+), Maastricht, The Netherlands; ³Department of Cardiac Surgery, Circolo Hospital, University of Insubria, Varese, Italy; ⁴Milano Bicocca University, Milan, Italy; ⁵Department for the Treatment and Study of Cardiothoracic Diseases and Cardiothoracic Transplantation, ⁶Department of Anesthesia and Intensive Care, IRCCS-ISMETT (Istituto Mediterraneo per i Trapianti e Terapie ad alta specializzazione), Palermo, Italy; ⁷De Gasperis Cardio Center, ASST Grande Ospedale Metropolitano Niguarda Ca' Granda, Milan, Italy; ⁸Department of Clinical Sciences and Community Health, Università degli Studi, Milan, Italy

Correspondence to: Paolo Meani, MD. Cardio-Thoracic Surgery Department, Maastricht University Medical Centre, P. Debyelaan 25 - 6202 AZ, Maastricht, The Netherlands. Email: paolo.meani@mumc.nl

Extracorporeal membrane oxygenation (ECMO) in the veno-arterial (VA) configuration is an established method for the treatment of refractory cardiogenic shock. Such a condition characterizes the postoperative course of approximately 1% of cardiac surgery patients. Although some studies have reported ECMO-related short-term results, little is known about the long-term outcomes of VA-ECMO therapy in the post-cardiotomy setting. Therefore, an extensive literature search was conducted regarding articles published after 1990 reporting postoperative ECMO use. PubMed, EMBASE and Web of Science were searched for sources. In-hospital mortality was high in post-cardiotomy VA-ECMO patients, ranging from 24.8% to 52%. Long-term results were poorly reported. However, based on the limited information available, hospital survivors showed a favorable outcome, with improvement in overall clinical condition, quality of life and limited hospital readmission for cardiac-related events. To conclude, in-hospital outcome in post-cardiotomy ECMO is often unfavorable, post-discharge results show satisfactory condition, with stable improvement of overall patient clinical status and low rate of hospital readmission and cardiac-related adverse events. Data reporting is, however, scarce and hence new and detailed studies are still warranted to investigate such aspects.

Keywords: Extracorporeal membrane oxygenation (ECMO); cardiogenic shock; heart failure; cardiac surgery



Submitted Nov 07, 2018. Accepted for publication Dec 01, 2018.

doi: 10.21037/acs.2018.12.04

View this article at: <http://dx.doi.org/10.21037/acs.2018.12.04>

Introduction

Since the first implantation by Hill and colleagues (1), extracorporeal membrane oxygenation (ECMO) has become an established treatment option for adult patients with refractory cardiogenic shock (2), providing temporary cardiac and respiratory support. Approximately 1% of

adult patients who undergo cardiac surgery develop post-cardiotomy cardiogenic shock (PCS) and require mechanical circulatory support (MCS) beyond the conventional medical or mechanical treatments to support the failing heart (1-5). Because of its quick availability, easy and rapid applicability and reliability, ECMO support has become a more and more attractive option, particularly in post-

cardiotomy (PC) setting (1-3). Another option available is short-term ventricular assist devices (VADs), either uni- or biventricular, but complexity, lack of lung support, and costs has limited its wide application (6-8).

PC-ECMO has become the most frequent indication for ECMO in the United States (3), however, its exponential increase has not been paralleled by improved results, but rather to the contrary (5). Although during the last 20 years advances in technique and management were achieved, ECMO results have not been satisfactory. Prevalence, trends of use, indications and implant information, types of patient, specific procedures for whom it is being used, in-hospital patient management and short- and long-term outcomes have peculiar features in the PC-ECMO setting. Information on post-hospital discharge outcomes after veno-arterial (VA)-ECMO (ranging from 1 year to 10 years) have shown overall survival rates ranging from 17% to 40% (9-11), but little attention has been given to other factors related to the long-term prognosis (12-14). Furthermore, despite numerous publications that have shown ECMO-related short and medium-term results, specific reports about PC-ECMO have been limited and usually related to single-center experience. In addition, some reports have shown some critical illness in survivors, with some long-term functional impairment in physical, cognitive and psycho-social settings (15,16). A targeted analysis of these aspects would have a relevant implication in the daily practice of ECMO users. This review, therefore, had the aim to present the current information about PC-ECMO in adult patients, providing an evaluation of the long-term mortality and morbidity for VA-ECMO.

Methods

A literature search was run on PubMed, EMBASE and Web of Science covering the time period between January 1990 and August 2018. Publications that appeared before 1990 were not included due to the remarkable dissimilarity in intensive care unit management, surgical techniques and ECMO-related technology. Two operators independently performed the computerized literature search using the following keywords: “Postoperative extracorporeal membrane oxygenation”, “Postoperative extracorporeal life support”, “Post-Cardiotomy”, “Mechanical Circulatory Support”, “Temporary Mechanical Circulatory Support”, “Perioperative Cardio-Pulmonary Assistance”, “Perioperative Cardio-Pulmonary Failure”, “Percutaneous Mechanical Circulatory Support”, “Postoperative

Cardiogenic Shock”, “Postoperative Cardiac Arrest”, “Post-cardiotomy Cardiac Arrest”, “Post-Cardiotomy Cardiac Failure”, “Post-Cardiotomy Complications”, “ARDS and cardiac surgery”, “Post-Cardiotomy Respiratory Dysfunction”. Information regarding target populations, ECMO implantation site and major outcomes were collected and systematically recorded. Only publications in English were considered.

Only studies specifically conducting on ECMO in peri- or post-operative adults (>18 years of age) undergoing cardiac surgery (see *Table 1*) were collected, leading to a careful assessment of short-term outcomes and evaluation of post-discharge follow-up. Exclusion criteria included published PC-ECMO series, case reports, or case series, reporting less than 10 patients, as these would not allow for a meaningful cohort analysis of PC-ECMO patients.

Results

Overall, 242 published papers were selected, reviewed and carefully analyzed. The flow-chart showing the literature search and selection for final information analysis is shown in *Figure 1*. *Table 1* shows the main results. Fifteen studies were selected, included 2,852 patients supported with VA-ECMO due to PCS. The mean age was 63.7 years. VA-ECMO was placed using central and peripheral cannulation. In-hospital survival ranged from 24.8% to 52%, whereas 1-year survival ranged from 16.5% to 47.6%. Regarding long-term survival, the best rate reported was 37% (*Table 1*). However, each study has considered different timing, therefore any consistent and comprehensive evaluation was not feasible. A total of six articles, enrolling 2,033 patients, reported 5-year survival, while only two studies (1,188 subjects) provided data for 10-year survival. Corresponding reported survival ranged from 13.7% to 32% and from 9.7% to 17.6%, respectively. The created variable regarding the survival difference, representing the difference between one-year and in-hospital survival, ranged from 0% to 21%, thereby confirming that long-term mortality was relatively low in these patients, although these findings were hampered by the maximum follow-up at 10 years from the original operation.

Table 2 shows the described major outcomes found in this literature selection. These mainly consisted of five fields reported: New York Heart Association Class, re-hospitalizations, medical expenditures, quality of life and specific outcomes regarding dead and survived patients. From the limited data and information available, it is

Table 1 Postcardiotomy ECMO studies

Author reference	Year of publication	Patient	Mean age (years)	In-hospital survival (%)	1-year survival (%)	Long-term survival (years) (%)
Magovern (17)	1994	21	61.6±2.2	52	47.6	NA
Kodera (18)	1996	17	41.4	47.1	NA	5 (30.9)
Ko (19)	2002	76	56.8±15.9	26.3	45.4	2.7 (23.0)
Doll (20)	2004	219	61.3±12.1	24	NA	5 (16.8)
Zhang (21)	2006	32	55.4±11.9	25	NA	3.8 (12.5)
Bakhtiary (22)	2008	45	60.1±13.6	29	29	3 (22.0)
Hsu (23)	2010	51	63±15.7	24.8	16.5	NA
Rastan (24)	2010	517	63.5±11.2	30.4	NA	5 (13.7)
Unosawa (25)	2013	47	64.4±12.5	40.3	24.5	3 (27.7) 5 (20.1) 10 (17.6)
Saxena (26)	2015	45	76.8±4.6	24.4	24.4	3 (17.7)
Khorsandi (27)	2016	16	71	51	29.8	2 (31.0)
Distelmaier (28)	2016	385	65	32	NA	3.6 (32.0)
Chen (29)	2017	1,141	63.8±13.2	38.3	24.1	5 (17.7) 10 (9.7)
Guihaire (30)	2017	92	64.5	42	39	2 (37.0) 5 (32.0)
Biancari (31)	2017	148	NA	33.8	31	2 (27.9) 3 (26.1)
Total/range	–	2,852	63.7	24.8–52	16.5–47.6	2–10 (9.7–37)

ECMO, extracorporeal membrane oxygenation.


**Figure 1** Literature search and selection flow chart.

Table 2 Postcardiotomy ECMO major outcomes

Outcomes	Details/results
NYHA class	
Ko (19)	NYHA class II or I
Doll (20)	NYHA class II
Bakhtiary (22)	NYHA class II or III
Unosawa (25)	NYHA class I or II
Saxena (26)	NYHA class I, II, III and IV
Khorsandi (27)	NYHA class I-II
Readmission	
Doll (20)	Rate: 37.8% Causes: routine cardiac investigations, pneumonia, non-cardiac-related surgical interventions
Bakhtiary (22)	Rate: 60% Causes: pneumonia, end-stage renal failure requiring dialysis
Chen (29)	Mean N of readmissions: 0.68; 95% CI, 0.60–0.78 Higher in PC patients support with ECMO
Expenditures	
Chen (29)	Greater in PC patients support with ECMO
Quality of life	
Guihaire (30)	Self-care without any limitations in all cases, minor problems considering other dimensions
Specific outcomes	
Ko (19)	Sudden death, pneumonia
Hsu (23)	Cardiac arrest, severe infection
Unosawa (25)	Cardiac events, heat stroke, ruptured iliac artery aneurysm, subarachnoid hemorrhage, pneumonia unknown cause
Saxena (26)	End-stage heart disease
Guihaire (30)	Major cardiac event, non-cardiac events
Bakhtiary (22)	Congestive heart failure, pneumonia, end-stage renal failure
NYHA, New York Heart Association; PC, post-cardiotomy; N, number; CI, confidence interval; ECMO, extracorporeal membrane oxygenation.	

evident that overall PC-ECMO patients have a favorable prognosis, with the vast majority maintaining a satisfactory clinical condition, limited readmission rate and favorable quality of life.

Discussion

The incidence of refractory PCS in adult cardiac surgical patients ranges from 0.5% to 1.5% (24,31,32). PCS is a rare but dreadful complication. The first PCS treatment step is pharmacologic. However, the overwhelming majority of PCS is unresponsive to high dosages of inotropes. Therefore, intra-aortic balloon pump support is frequently adopted, even if this strategy presents well-known limitations, particularly during active resuscitation (33). As a direct consequence, VA-ECMO has been increasingly considered and applied in PCS. Such a temporary MCS is a salvage therapy, often representing the only prompt and life-saving technique in refractory PCS cases. Although VA-ECMO support is a well-established and effective procedure in the presence of PCS, the in-hospital mortality rate is still high, with an unsatisfactory in-hospital outcome. In fact, the reported in-hospital survival has ranged from 24.8% to 52% in several case series, a 1-year survival ranging from 16.5% to 47.6% and gradually ranging from almost 10% to 37% when the late survival is considered (20-30). These data clearly show how the majority of PC-ECMO patients die during the hospitalization, showing a fall in the mortality rate after being discharged. This would mean that achieving the discharge might be one of the most important predictors of good long-term outcome. Therefore, it is clear that high in-hospital mortality remains the Achilles' heel of ECMO. The importance of survival to discharge—and not just successful weaning—was illustrated by Ko *et al.*, who studied 76 patients undergoing ECMO support for refractory PCS. Although 60.5% of this cohort was successfully weaned from ECMO, only 26.3% survived to discharge. However, all survivors showed New York Heart Association (NYHA) I and II functional class at 32±22 months of follow-up (19).

However, the current literature doesn't allow to make any substantial consideration regarding long-term outcomes. Lack of data and absence of standardized data collection made our analysis extremely undetailed and difficult to interpret. Under this light, the overall literature clearly shows a lot of cloudy aspects. von Bahr *et al.* reported a high risk of death, 15% within 90 days of hospital discharge increasing to 24% at long-term follow-up of 3 years (34).

In contrast, de Waha *et al.* showed 15.7% survival with satisfactory functional outcomes at 18 months in patients with refractory cardiogenic shock undergoing ECLS (35). Therefore, another crucial point that should be detailed better is the quality of life of survivors.

Other outcomes

The NYHA class is a remarkable scale for defining patient functional status (36). Six out of sixteen papers reported this parameter as part of long-term results in PC-ECMO patients. Doll *et al.* showed all survivors in NYHA class II at late follow-up (20). On the contrary, the experience presented by Saxena *et al.* (26), interestingly described a heterogeneous functional status, which included patients in class III and IV. However, the overwhelming majority of these studies reported survivors in NYHA class I or II (20-30). This might lead to speculation that the PC-ECMO survivors, once overcoming the ECMO-related hospital course, may show a favorable long-term functional status. Guihaire *et al.* supported the latter finding by measuring the long-term quality of life (30). In this study, the health-related quality of life was assessed among PC-ECMO survivors 2 years after ECMO support using the EQ-5D survey (37) and the EQ visual analogue scale. No self-care limitations were found in any cases. Minor problems, however, were found considering other dimensions came out (e.g., mobility, usual activities, pain and anxiety) (37). Therefore, PC-ECMO survivors seem to achieve a long-term good functional status and quality of life. On the other hand, three papers included readmission rate in their analysis. The latter fluctuated from 37.8% to 60%. Under this light, Chen *et al.* (29) observed a greater mean number of readmissions for any cause during the follow-up in patients who required ECMO compared with those who didn't need the ECMO support after surgery. As a consequence, patients supported with ECMO for PCS might have an increased need of future hospitalizations. Chen *et al.* also found greater medical expenditure in PC-ECMO group during only the first year, which disappeared in the following checks. This finding might be related with the increase number of readmissions in this population, underlying the high disease burden amongst these patients. Finally, scant data are available regarding the specific causes of death or the new onset of morbidities in survivors. Nevertheless, infections and cardiac events might be considered the most common causes of death or morbidity (20-30).

Conclusions

In conclusion, the long-term survival in PC-ECMO seems to be strongly influenced by the early hospitalized critical phase, after which the survival slightly drops but remains satisfactory. Little is known, however, regarding other long-term outcome parameters after ECMO treatment in PC setting. The current lack of dedicated registries, limited data collection within studies and post-discharge follow-up in PC-ECMO patients, and a few centers with this specific experience and high number of cases, are likely the most important causes for such poor information. Further investigations are urgently needed in order to better show post-discharge outcomes in adult patients submitted to PC-ECMO.

Acknowledgements

None.

Footnote

Conflicts of Interest: Prof. Lorusso declares that he is Principal Investigator of the PERSIST-AVR Study (sponsored by LivaNova) and is Consultant for Medtronic. The other authors have no conflicts of interest to declare.

References

1. Hill JD, O'Brien TG, Murray JJ, et al. Prolonged extracorporeal oxygenation of acute posttraumatic respiratory failure (shock-lung syndrome). *N Engl J Med* 1972;286:629-34.
2. McCarthy FH, McDermott KM, Kini V, et al. Trends in U.S. extracorporeal membrane oxygenation use and outcome: 2002-2012. *Semin Thorac Cardiovasc Surg* 2015;27:81-8.
3. Sauer CM, Yuh DD, Bonde P. Extracorporeal membrane oxygenation use increased 433% in adults in the United States from 2006 to 2011. *ASAIO J* 2015;61:31-6.
4. Maxwell BG, Powers AJ, Sheikh AY, et al. Resource use trends in extracorporeal membrane oxygenation in adults: an analysis of the Nationwide Inpatient Sample 1998-2009. *J Thorac Cardiovasc Surg* 2014;148:416-21.e1.
5. Whitman GJR. Extracorporeal membrane oxygenation for the treatment of postcardiotomy shock. *J Thorac Cardiovasc Surg* 2017;153:95-101.
6. Goldstein DJ, Oz MC. Mechanical support for

- postcardiotomy cardiogenic shock. *Semin Thorac Cardiovasc Surg* 2000;12:220-8.
7. Curtis JJ, McKenney-Knox CA, Wagner-Mann CC. Postcardiotomy centrifugal assist: a single surgeon's experience. *Artif Organs* 2002;26:994-7.
 8. Jurmann MJ, Siniawski H, Erb M, et al. Initial experience with miniature axial flow ventricular assist devices for postcardiotomy heart failure. *Ann Thorac Surg* 2004;77:1642-7.
 9. Combes A, Leprince P, Luyt CE, et al. Outcomes and long-term quality-of-life of patients supported by extracorporeal membrane oxygenation for refractory cardiogenic shock. *Crit Care Med* 2008;36:1404-11.
 10. Lee SH, Chung CH, Lee JW, et al. Factors predicting early- and long-term survival in patients undergoing extracorporeal membrane oxygenation (ECMO). *J Card Surg* 2012;27:255-63.
 11. Winters BD, Eberlein M, Leung J, et al. Long-term mortality and quality of life in sepsis: a systematic review. *Crit Care Med* 2010;38:1276-83.
 12. Bottiroli M, Maraffi T, Decaria D, et al. In-hospital and long-term mortality after venoarterial ECMO for refractory cardiogenic shock. *Crit Care* 2015;19:P281.
 13. Hodgson CL, Hayes K, Everard T, et al. Long-term quality of life in patients with acute respiratory distress syndrome requiring extracorporeal membrane oxygenation for refractory hypoxaemia. *Crit Care* 2012;16:R202.
 14. Lindén VB, Lidégran MK, Frisen G, et al. ECMO in ARDS: a long-term follow-up study regarding pulmonary morphology and function and health-related quality of life. *Acta Anaesthesiol Scand* 2009;53:489-95.
 15. Hashem MD, Nallagangula A, Nalamalapu S, et al. Patient outcomes after critical illness: a systematic review of qualitative studies following hospital discharge. *Crit Care* 2016;20:345.
 16. Herridge MS, Tansey CM, Matte A, et al. Functional disability 5 years after acute respiratory distress syndrome. *N Engl J Med* 2011;364:1293-304.
 17. Magovern GJ Jr, Magovern JA, Benckart DH, et al. Extracorporeal membrane oxygenation: preliminary results in patients with postcardiotomy cardiogenic shock. *Ann Thorac Surg* 1994;57:1462-8; discussion 1469-71.
 18. Kodera K, Kitamura M, Hachida M, et al. Biventricular bypass with oxygenation for postcardiotomy ventricular failure. *Artif Organs* 1996;20:724-7.
 19. Ko WJ, Lin CY, Chen RJ, et al. Extracorporeal membrane oxygenation support for adult postcardiotomy cardiogenic shock. *Ann Thorac Surg* 2002;73:538-45.
 20. Doll N, Kiaii B, Borger M, et al. Five-year results of 219 consecutive patients treated with extracorporeal membrane oxygenation for refractory postoperative cardiogenic shock. *Ann Thorac Surg* 2004;77:151-7.
 21. Zhang R, Kofidis T, Kamiya H, et al. Creatine kinase isoenzyme MB relative index as predictor of mortality on extracorporeal membrane oxygenation support for postcardiotomy cardiogenic shock in adult patients. *Eur J Cardiothorac Surg* 2006;30:617-20.
 22. Bakhtiari F, Keller H, Dogan S, et al. Venoarterial extracorporeal membrane oxygenation for treatment of cardiogenic shock: clinical experiences in 45 adult patients. *J Thorac Cardiovasc Surg* 2008;135:382-8.
 23. Hsu PS, Chen JL, Hong GJ, et al. Extracorporeal membrane oxygenation for refractory cardiogenic shock after cardiac surgery: predictors of early mortality and outcome from 51 patients. *Eur J Cardiothorac Surg* 2010;37:328-33.
 24. Rastan AJ, Dege A, Mohr M, et al. Early and late outcome of 517 consecutive patients treated with extracorporeal membrane oxygenation for refractory postcardiotomy cardiogenic shock. *J Thorac Cardiovasc Surg* 2010;139:302-11.
 25. Unosawa S, Sezai A, Hata M, et al. Long-term outcomes of patients undergoing ECMO for refractory postcardiotomy cardiogenic shock. *Surg Today* 2013;43:264-70.
 26. Saxena P, Neal J, Joyce LD, et al. Extracorporeal membrane oxygenation support in postcardiotomy elderly patients: the Mayo Clinic experience. *Ann Thorac Surg* 2015;99:2053-60.
 27. Khorsandi M, Shaikhrezai K, Prasad S, et al. Advanced mechanical circulatory support for post-cardiotomy cardiogenic shock: a 20-year outcome analysis in a non-transplant unit. *J Cardiothorac Surg* 2016;11:29-35.
 28. Distelmaier K, Schrutka L, Binder C, et al. Cardiac arrest does not affect survival in post-operative cardiovascular surgery patients undergoing extracorporeal membrane oxygenation. *Resuscitation* 2016;104:24-7.
 29. Chen SW, Tsai FC, Lin YS, et al. Long-term outcomes of extracorporeal membrane oxygenation support for postcardiotomy shock. *J Thorac Cardiovasc Surg* 2017;154:469-77.e2.
 30. Guihaire J, Dang Van S, Rouze S, et al. Clinical outcomes in patients after extracorporeal membrane oxygenation support for post-cardiotomy cardiogenic shock: a single-centre experience. *Interact Cardiovasc Thorac Surg* 2017;25:363-9.
 31. Biancari F, Dalén M, Perrotti A. Venoarterial

- extracorporeal membrane oxygenation after coronary artery bypass grafting: Results of a multicenter study. *Int J Cardiol* 2017;241:109-14.
32. Fux T, Holm M, Corbascio M, et al. Venoarterial extracorporeal membrane oxygenation for postcardiotomy shock: risk factors for mortality. *J Thorac Cardiovasc Surg* 2018;156:1894-902.e3.
 33. Bozbuga N. Postcardiotomy extracorporeal membrane oxygenation for refractory cardiogenic shock. *J Thorac Cardiovasc Surg* 2018;156:1903-4.
 34. von Bahr V, Hultman J, Eksborg S, et al. Long-term survival in adults treated with extracorporeal membrane oxygenation for respiratory failure and sepsis. *Crit Care Med* 2017;45:164-70.
 35. de Waha S, Fuernau G, Desch S, et al. Long-term prognosis after extracorporeal life support in refractory cardiogenic shock: results from a real-world cohort. *EuroIntervention* 2016;11:1363-71.
 36. Holland R, Rechel B, Stepien K, et al. Patients' self-assessed functional status in heart failure by New York Heart Association class: a prognostic predictor of hospitalizations, quality of life and death. *J Card Fail* 2010;16:150-6.
 37. Rabin R, de Charro F. EQ-5D: a measure of health status from the EuroQol Group. *Ann Med* 2001;33:337-43.

Cite this article as: Meani P, Matteucci M, Jiritano F, Fina D, Panzeri F, Raffa GM, Kowalewski M, Morici N, Viola G, Sacco A, Oliva F, Alyousif A, Heuts S, Gilbers M, Schreurs R, Maessen J, Lorusso R. Long-term survival and major outcomes in post-cardiotomy extracorporeal membrane oxygenation for adult patients in cardiogenic shock. *Ann Cardiothorac Surg* 2019;8(1):116-122. doi: 10.21037/acs.2018.12.04