



doi: 10.21037/acs-2023-rcabg-0198

Cite this article as: Van Puyvelde J, Baudo M, Torregrossa G, Oosterlinck W. The role of robotic coronary artery bypass grafting in the current practice of surgical myocardial revascularization. *Ann Cardiothorac Surg* 2024. doi: 10.21037/acs-2023-rcabg-0198

This is a PDF file of an edited manuscript that has been accepted for publication. As a service to our customers we are providing this Online First version of the manuscript. The manuscript has undergone copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The role of robotic coronary artery bypass grafting in the current practice of surgical myocardial revascularization

Joeri Van Puyvelde^{1#}, Massimo Baudo^{1#}, Gianluca Torregrossa², Wouter Oosterlinck¹

¹Department of Cardiovascular Sciences, Research Unit of Cardiac Surgery, University Hospitals Leuven, KU Leuven, Leuven, Belgium;

²Department of Cardiothoracic Surgery, Lankenau Heart Institute, Lankenau Medical Center, Main Line Health, Wynnewood, PA, USA

[#]These authors contributed equally to this work as co-first authors.

Correspondence to: Joeri Van Puyvelde, MD. Department of Cardiovascular Sciences, Research Unit of Cardiac Surgery, University Hospitals Leuven, KU Leuven, Herestraat 49, 3000 Leuven, Belgium. Email: joeri.vanpuyvelde@uzleuven.be.

Keywords: Myocardial revascularization; robotic minimally invasive direct coronary artery bypass grafting (robotic MIDCAB); hybrid coronary revascularization; coronary artery disease

Submitted Nov 24, 2023. Accepted for publication Mar 05, 2024. Published online Apr 03, 2024.

doi: 10.21037/acs-2023-rcabg-0198

View this article at: <https://dx.doi.org/10.21037/acs-2023-rcabg-0198>

The field of coronary artery bypass grafting (CABG) has evolved significantly, giving rise to refined techniques and improved patient outcomes (1). The introduction of minimally invasive techniques, starting with minimally invasive direct coronary artery bypass grafting (MIDCAB) and advancing to robotic MIDCAB and totally endoscopic coronary artery bypass grafting (TECAB), has further revolutionized the field of cardiac surgery (2). These minimally invasive techniques address the limitations of traditional CABG by enabling surgery through small incisions without the use of cardiopulmonary bypass, offering reduced trauma, shorter hospital stays, and faster recovery. Furthermore, the integration of these minimally invasive methods with interventional techniques in hybrid coronary revascularization (HCR) strategies has amalgamated the benefits of both approaches (3).

Surgical revascularization using internal mammary arteries (IMAs) has demonstrated superior long-term patency and improved survival compared to percutaneous coronary intervention (PCI) in stable coronary artery disease (4). Robotic MIDCAB offers the potential for enhanced precision, dexterity, and visualization, facilitating more intricate anastomoses and potentially improving long-term graft patency, thereby contributing to improved patient outcomes (5).

Despite the potential advantages of robotic MIDCAB,

there have been several challenges that hinder its widespread adoption. These include the high initial and procedural cost of robotic systems, the need for specialized expertise, and the unjustly feared learning curve associated with these techniques. Nevertheless, research has indicated that outcomes can significantly improve when working with a well-trained team, following standardized procedures, and adopting the right approach (6). Despite these obstacles, the number of centers with robotic programs is steadily increasing (7).

Heart team discussions play a crucial role in evaluating individual patients and determining the most appropriate treatment approach. By facilitating meaningful interdisciplinary discussions, patients can benefit from the combined expertise of both specialties. The presence of a robotic MIDCAB program in surgical departments lays a solid foundation for collaborative hybrid revascularization procedures with cardiologists (8). Robotic MIDCAB presents an opportunity for cardiologists to assess patients who may not be suitable for traditional cardiac surgery for robotic MIDCAB or hybrid coronary revascularization. This allows for more comprehensive revascularization across a wider spectrum of patients. Specifically, robotic MIDCAB with IMA to left anterior descending artery (LAD) may offer advantages over PCI for isolated LAD lesions, particularly in younger individuals or cases requiring longer stents,

which pose a higher risk of restenosis (9). Additionally, robotic MIDCAB can complement PCI in addressing multivessel coronary artery disease, leading to personalized and optimal treatment strategies. HCR, especially when combining robotic MIDCAB with PCI, presents a superior approach in cases where certain vessels are not optimal targets for surgical revascularization (3). This combined strategy enables full revascularization by allowing for PCI of vessels that may pose challenges for traditional surgical intervention. By combining the accuracy and flexibility of PCI with the enduring patency of IMAs used in robotic MIDCAB, complete and durable revascularization can be achieved, potentially leading to improved outcomes. In the context of urgent non-ST-elevation or ST-elevation myocardial infarction cases, there exists the potential for additional benefit through a dual strategy involving immediate PCI to address the non-LAD culprit lesion during the ischemic event, followed by subsequent complete revascularization of the LAD lesion using robotic MIDCAB on a longer-term basis. This approach not only addresses the acute ischemic event promptly but also provides a pathway for comprehensive and durable revascularization through robotic MIDCAB, potentially contributing to improved long-term outcomes for these high-risk patients. Furthermore, HCR involving robotic MIDCAB provides an opportunity for optimal revascularization in patients with severe coronary artery disease, advanced age, or comorbidities that render them unsuitable for high-risk CABG surgery involving a median sternotomy. Patients at the highest risk of mortality with conventional CABG have shown considerable benefits from off-pump CABG due to the reduced physiological stress it imposes, such as less systemic inflammation and haemodilution, compared to CABG performed on cardiopulmonary bypass (10,11). Robotic MIDCAB, with its minimally invasive approach, further mitigates physiological stress and inflammation associated with surgery, providing additional benefits for these high-risk patients. This approach extends optimal care to the most vulnerable patients, including those awaiting transcatheter valve intervention, individuals with chronic kidney insufficiency and moderate coronary artery disease involving the proximal LAD, as well as morbidly obese patients at a heightened risk of postoperative sternotomy complications. The emphasis on rapid postoperative recovery is particularly significant in these cases, and minimally invasive/hybrid revascularization demonstrates substantial benefits in achieving swift extubation, reducing postoperative pain, shortening hospital stays, and facilitating

prompt rehabilitation.

In conclusion, robotic MIDCAB holds immense potential in the realm of HCR, offering a minimally invasive approach with the potential for improved patient outcomes. However, current guidelines lack specific recommendations for robotic MIDCAB and HCR, and differences exist between European and American guidelines. It is imperative to consider implementing recommendations for HCR with robotic MIDCAB in future guidelines to ensure standardized and evidence-based approaches to myocardial revascularization. As technological advancements progress and collaborative efforts between surgery and interventional medicine strengthen, the future of robotic MIDCAB in hybrid revascularization strategies appears promising.

Acknowledgments

Funding: None.

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

Open Access Statement: This is an Open Access article distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Head SJ, Milojevic M, Taggart DP, et al. Current Practice of State-of-the-Art Surgical Coronary Revascularization. *Circulation* 2017;136:1331-45.
2. Melly L, Torregrossa G, Lee T, et al. Fifty years of coronary artery bypass grafting. *J Thorac Dis* 2018;10:1960-7.
3. Van den Eynde J, Sá MP, De Groote S, et al. Hybrid coronary revascularization versus percutaneous coronary intervention: A systematic review and meta-analysis. *Int J Cardiol Heart Vasc* 2021;37:100916.
4. Doenst T, Haverich A, Serruys P, et al. PCI and CABG for Treating Stable Coronary Artery Disease: JACC Review

- Topic of the Week. *J Am Coll Cardiol* 2019;73:964-76.
5. Bonatti J, Wallner S, Crailsheim I, et al. Minimally invasive and robotic coronary artery bypass grafting—a 25-year review. *J Thorac Dis* 2021;13:1922-44.
 6. Patrick WL, Iyengar A, Han JJ, et al. The learning curve of robotic coronary arterial bypass surgery: A report from the STS database. *J Card Surg* 2021;36:4178-86.
 7. Cerny S, Oosterlinck W, Onan B, et al. Robotic Cardiac Surgery in Europe: Status 2020. *Front Cardiovasc Med* 2021;8:827515.
 8. Van den Eynde J, Bennett J, McCutcheon K, et al. Heart team 2.0: A decision tree for minimally invasive and hybrid myocardial revascularization. *Trends Cardiovasc Med* 2021;31:382-91.
 9. Diegeler A, Thiele H, Falk V, et al. Comparison of stenting with minimally invasive bypass surgery for stenosis of the left anterior descending coronary artery. *N Engl J Med* 2002;347:561-6.
 10. Biglioli P, Cannata A, Alamanni F, et al. Biological effects of off-pump vs. on-pump coronary artery surgery: focus on inflammation, hemostasis and oxidative stress. *Eur J Cardiothorac Surg* 2003;24:260-9.
 11. Puskas JD, Thourani VH, Kilgo P, et al. Off-pump coronary artery bypass disproportionately benefits high-risk patients. *Ann Thorac Surg* 2009;88:1142-7.

Cite this article as: Van Puyvelde J, Baudo M, Torregrossa G, Oosterlinck W. The role of robotic coronary artery bypass grafting in the current practice of surgical myocardial revascularization. *Ann Cardiothorac Surg* 2024. doi: 10.21037/acs-2023-rcabg-0198

